

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
8 April 2004 (08.04.2004)

PCT

(10) International Publication Number
WO 2004/028498 A1

(51) International Patent Classification⁷: **A61K 7/16**

(21) International Application Number:
PCT/GB2003/004162

(22) International Filing Date:
23 September 2003 (23.09.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0222091.1 24 September 2002 (24.09.2002) GB

(71) Applicants (*for all designated States except US*): **THE BOOTS COMPANY PLC** [GB/GB]; 1 Thane Road West, Nottingham NG2 3AA (GB). **THE UNIVERSITY OF SHEFFIELD** [GB/GB]; Firth Court, Western Bank, Sheffield, South Yorkshire S10 2TE (GB).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **GUAN, Yue, Hugh** [GB/GB]; University of Sheffield, School of Dentistry, Claremont Crescent, Sheffield, South Yorkshire S10 2TA (GB). **LILLEY, Terence, Henry** [GB/GB]; 451 Greystones Road, Sheffield, South Yorkshire S11 7BY (GB).

(74) Agent: **JONES, Stephen, Anthony**; Adamson Jones, Broadway Business Centre, 32a Stoney Street, Nottingham NG1 1LL (GB).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

— *of inventorship (Rule 4.17(iv)) for US only*

Published:

— *with international search report*
— *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **DENTAL COMPOSITIONS AND METHODS**

(57) Abstract: A dental composition comprises a fluorescent whitening agent. The fluorescent whitening agent is most preferably 4,4'-bis(2-sulfostyryl)biphenyl or a salt or other soluble derivative thereof, eg the disodium salt. The composition may also comprise an additional tooth whitening agent. Methods of whitening the teeth are also described.

WO 2004/028498 A1

Title – Dental Compositions and Methods

This invention relates to dental compositions and methods, and in particular to compositions and methods for enhancing or improving the whiteness of teeth.

5

Staining or discolouration of teeth can arise as a result of a number of factors, including

- 1) exposure of the teeth to staining agents such as tea, coffee, smoke etc, which can cause entrapment or attachment of stain molecules to the tooth surface;
- 10 2) ingestion of certain materials, eg certain drugs such as tetracycline, during tooth formation, which can cause discolouration of the tooth enamel;
- 3) trauma to the tooth that can cause leakage of blood breakdown products into the enamel and thus discolouration.

15 A number of different methods for tooth whitening are known. These include:

- 1) Physical abrading of the stain to remove it. This involves the use of harsher abrasives than are found in typical tooth-whitening toothpaste formulations. Brushing or a similar polishing or scrubbing action is required to complement the action. This approach suffers from the disadvantage that it may cause excessive
 20 abrasion to the tooth enamel.
- 2) Oxidising or bleaching agents such as peroxides are widely used to effect removal of both extrinsic and intrinsic stains. Such agents can be applied either as pastes or gels. However, the treatment may require considerable time to achieve good results. The concentration of oxidising agent may be increased to
 25 reduce the time required to achieve effective results, but this may lead to safety concerns.
- 3) Enzymatic systems comprising proteases such as papain that are claimed to remove stain by dissolution of the pellicle entrapping the stain.
- 4) Chelating agents have also been used to whiten teeth by interacting either
 30 with inorganic components of the tooth and tartar or with stain molecules. Typical examples of such agents are pyrophosphates and polyphosphates.

The natural colour of the tooth is very rarely pure white, mainly because the dentin, which contains a substantial amount of collagen, is yellowish and the enamel above it is not completely opaque. The "whitening" approaches 3) and 4) described above are primarily based on interactions with acquired enamel pellicle, plaque and dental calculus. The mechanism employed is to either remove or weaken these tooth surface adsorbents and precipitates. Approach 1) simply polishes the enamel using abrasives and thus removes stains. However, selection of abrasives always involves a balance between its efficacy and mechanical destruction of the enamel. The use of bleaching agents, as in approach 2), involves chemical removal of the colour of organic molecules on or within the tooth surface and thus achieves a better cosmetic whitening effect.

Fluorescent whitening agents have been known for many years. Most molecules categorised as fluorescent whitening agents contain chromophores which contain conjugated double bonds that absorb UV light. By re-emitting the absorbed light at higher wavelength regions, within the visible spectrum, the fluorescent whitening agents can alter the visual appearance of a substrate to which they are applied. The optical properties of the chromophores can be varied by electron donors or acceptors. Hitherto, however, the use of such compounds has been confined to industrial fields such as the textile, paper and detergent industries.

It has now been surprisingly found that fluorescent whitening agents can advantageously be incorporated in dental compositions, and can be used in methods for improving or enhancing the whiteness of teeth. The present invention thus provides dental compositions and methods that overcome or substantially mitigate the disadvantages of the prior art approaches described above.

According to a first aspect of the invention, there is provided a dental composition comprising a fluorescent whitening agent and a dentally acceptable diluent or carrier.

The composition according to the invention is advantageous primarily in that it is found that the composition, containing the fluorescent whitening agent, is effective in improving the whiteness of teeth to which it is applied. Such an effect is believed to be due to an adherence of the fluorescent whitening agent to the teeth, presumably to the pellicle, the outer layer of proteins and carbohydrates that coats the tooth enamel. Such adherence, if indeed it does occur, could not have been predicted from the known industrial applications of fluorescent whitening agents.


As used herein, a "dental composition" is any composition suitable and intended for administration to the buccal cavity for the purpose of improving the appearance of the teeth. In the context of this invention, this means improving or enhancing the whiteness of the teeth. The dental composition may, for instance, be formulated as a toothpaste, mouthrinse, toothgel, tooth paint or a dental gel.

By a "fluorescent whitening agent" is meant a compound that absorbs invisible ultra-violet light and which re-emits the light in the visible spectral range. The compound thus absorbs light of wavelength less than, say, 380nm, and re-emits light in the wavelength range 380nm(violet) to 780nm (red). To achieve a whitening effect, the re-emitted light should complement the light reflected from the substrate to which the fluorescent whitening agent is applied, so that a more uniform spectral distribution is achieved across the whole visible range. It is particularly preferred, in order to compensate for the yellow cast of discoloured teeth, that the fluorescent whitening agent should emit in the blue region of the visible spectrum, eg in the range 400nm to 450nm.

The fluorescent whitening agent may be any compound that exhibits appropriate spectroscopic properties, and which is acceptable for administration in the manner contemplated by this invention, eg from the point of view of its toxicological properties.


Fluorescent whitening agents of potential utility in the present invention include the following classes of compound:

a) bis-styrylbiphenyl compounds, ie compounds having the following chromophore system:




5


b) derivatives of stilbene, ie compounds having the following chromophore system:



10 and in particular bis-triazineamine derivatives of such compounds having the following chemical backbone:




15 c) coumarins having the following chromophore system:




d) naphthalimide derivatives having the following chromophore system:

20



e) benzoxazole derivatives having the following chromophore system:




in which R represents a conjugated group, eg an ethylene group or a stilbene moiety (see above);

10 f) pyrazoline and bis-pyrazoline derivatives having the following chromophore systems:



15 It will be appreciated that all of the chromophore systems defined above may carry one or more substituent atoms or groups, and a great number of derivatives can be made by chemical substitution.

Preferred bis-styrylbiphenyl compounds of utility in the invention may be
20 represented by the general formula:




in which R_1 is $-\text{SO}_3\text{M}$ and R_2 , R_3 and R_4 , which may be the same or different, are selected from R_5 , $-\text{SO}_3\text{M}$, halogen (particularly Cl), $-\text{CN}$, $-\text{OC}(=\text{O})\text{R}_5$, $-\text{COOR}_5$, $-\text{SO}_2\text{N}(\text{R}_5)_2$ and $-\text{CON}(\text{R}_5)_2$, wherein R_5 represents hydrogen or C_{1-8} alkyl and M represents hydrogen or a Group I metal, eg Na, K or Li. Particularly preferred compounds are those in which R_3 is the same as R_1 , and R_2 and R_4 are the same and are selected from R_5 , halogen, $-\text{CN}$, $-\text{OC}(=\text{O})\text{R}_5$, $-\text{COOR}_5$, $-\text{SO}_2\text{N}(\text{R}_5)_2$ and $-\text{CON}(\text{R}_5)_2$.


Examples of bis-styrylbiphenyl compounds that may be used are 4,4'-bis(2-sulfo-4-chlorostyryl)biphenyl (sold under the name of Tinopal CBS) and disodium 4,4'-bis(3-sulfo-4-chlorostyryl)biphenyl, both of which are supplied by Ciba Speciality Chemicals plc.

Examples of bistriazineamino derivatives of stilbene are disodium 4,4'-bis[(4-anilino-6-morpholino-1,3,5-triazin-2-yl)amino]stilbene-2,2'-disulfonate (sold under the trade name Tinopal DMS), disodium 4,4'-bis[(4-anilino-6-(N-methyl-N-2-hydroxyethyl)amino-1,3,5-triazin-2-yl) amino]stilbene-2,2'-disulfonate, and disodium 4,4'-bis[(4-anilino-6-methylamino-1,3,5-triazin-2-yl)amino]stilbene-2,2'-disulfonate (all of which are supplied by Ciba Specialty Chemicals plc).

An example of a coumarin fluorescent whitening agent is Coumarin 102, the structure of which is:



An example of a naphthalimide fluorescent whitening agent is 1,8-naphthalimide, the structure of which is:



- 5 A particularly preferred fluorescent whitening agent is 4,4'-bis(2-sulfo-4-vinylphenyl)biphenyl or a salt or other soluble derivative thereof, eg the disodium salt, one such compound being sold under the trade name Tinopal CBS-X by Ciba Speciality Chemicals.
- 10 The invention may involve the use of a compound known for use as a fluorescent whitening agent in other fields of application (eg for application to textile or for use in a detergent composition), or the use of a compound having the required optical properties which is either a novel compound or a compound which, while known *per se*, has not previously been used as a fluorescent whitening agent.
- 15 The content of fluorescent whitening agents in the composition according to the invention will depend on a number of factors, including the particular fluorescent whitening agent used and the magnitude of the desired whitening effect. A distinction may also be drawn between compositions intended for clinical use by a
- 20 dental surgeon, and those for self-administration (eg at home) by the patient. In the former case, the concentration of fluorescent whitening agent may be up to 1000ppm, but is more commonly 500ppm or less, say between 50ppm and 500ppm. In the latter case, the concentration of fluorescent whitening agent will generally be much lower, say up to 100ppm, and more usually up to 50ppm, eg
- 25 between 5ppm and 50ppm.

According to another aspect of the invention, there is provided a method for improving or enhancing the appearance of the teeth, in particular for whitening of

the teeth, which method comprises application to the teeth of a dental composition comprising a fluorescent whitening agent and a dentally acceptable diluent or carrier.

- 5 To achieve maximal results, the treatment will normally be repeated at intervals for a period of several weeks or months. For instance, the treatment may be applied initially by a dental surgeon, who may also provide one or more subsequent treatments at intervals of, say, one or two days, or one or two weeks. Between these clinical applications of the composition, the patient may himself apply regular
- 10 treatments, eg once or several times each day.

In preferred embodiments, the method according to the invention involves treatment of the teeth, in addition to treatment with the fluorescent whitening agent, with one or more additional tooth whitening agents. Such other agents

15 may, for instance, be

- a) abrasive agents effective in physically removing stains from the tooth enamel;
- b) oxidising or bleaching agents such as peroxides;
- c) enzymatic systems, eg those including proteases such as papain; and
- 20 d) chelating agents, eg pyrophosphates and polyphosphates.

Thus, according to more specific aspects of the invention, there are provided

- a) a method for improving or enhancing the appearance of the teeth, in particular for whitening of the teeth, which method comprises application to the
- 25 teeth of an additional tooth whitening agent and a fluorescent whitening agent; and
- b) a dental composition comprising an additional tooth whitening agent and a fluorescent whitening agent, together with a dentally acceptable diluent or carrier.

The effects of the additional whitening agent and the fluorescent whitening agent

30 may be more than simply cumulative. It is believed that, at least in some instances, treatment of the teeth with the additional whitening agent may enhance the binding to the teeth of the fluorescent whitening agent, and thereby lead to a

greater-than-additive overall effect of the two agents. In addition, treatment with the additional whitening agent may lead to greater uniformity of the enhanced whiteness achieved by application of the fluorescent whitening agent.

- 5 The additional whitening agent may be any agent effective to improve the whiteness of the teeth. The preferred form of additional whitening agent, however, is an oxidising or bleaching agent.

10 Suitable forms of bleaching agent include compounds containing (or compounds capable, in use, of generating) peroxide or chlorite ions. Examples of peroxide or peroxide-generating compounds include hydrogen peroxide, sodium percarbonate, sodium perborate, sodium persilicate, sodium perphosphate, calcium peroxide and sodium peroxide. The presently most preferred peroxide bleaching agents are hydrogen peroxide or adducts such as urea peroxide (carbamide peroxide).

15 Examples of chlorite bleaching agents include alkali metal chlorites such as lithium chlorite, potassium chlorite and, especially, sodium chlorite, and alkaline earth metal chlorites such as calcium chlorite, magnesium chlorite and barium chlorite. Chlorite salts of other metals, and other salts, eg polymeric salts, may also be
20 used. Water-soluble chlorites are generally preferred. Such compounds may generate chlorine dioxide upon contact with an acid.

Particularly preferred bleaching agents for use in the invention are hydrogen peroxide and urea peroxide (carbamide peroxide).

25 The concentration of additional whitening agent used in the method according to the invention will depend on factors such as the nature of the additional whitening agent and the magnitude of the desired whitening effect. Again, a distinction can be drawn between clinical and home use. In the case of the preferred bleaching
30 agent hydrogen peroxide, the bleaching agent will typically be present in the composition at a level sufficient to generate a concentration of peroxide of up to 50% by weight in the case of a composition for clinical use, more typically up to

30%, and usually between 1% and 15%, typically from 5% to 15%. In compositions for home use, on the other hand, the concentration of bleaching agent will generally be much lower, usually less than 10%, and often less than 1%, eg 0.1%.

5

Application of the fluorescent whitening agent and the additional whitening agent may be simultaneous, in which case the fluorescent whitening agent and the additional whitening agent may be formulated in the same composition.

10 Alternatively, the additional whitening agent and the fluorescent whitening agent may be applied sequentially, in which case they will be formulated in separate compositions. In such a case, the additional whitening agent is applied first, followed by the fluorescent whitening agent.

15 The method according to the invention may be a clinical method carried out by, or under the direct supervision of a professional dental surgeon. Alternatively, the method may be carried out by a user at home, the composition of the invention being self-administered. The method may involve both types of treatment, these being an initial professional treatment, and optionally one or more subsequent
20 professional treatments, and subsequent maintenance treatment performed by the user.

In a presently favoured treatment regime, the fluorescent whitening agent and the additional tooth whitening agent are applied separately in a first stage of the tooth
25 whitening process, and then applied together in a second stage of the process. The first stage may be carried out by a dental surgeon, and the second stage may be carried out by the patient.

Application of the compositions used in the invention to the teeth may involve
30 simply bringing the composition into contact with the teeth, eg by introducing the composition into the buccal cavity and allowing it to disperse within that cavity.

Toothguards or the like may be used to constrain the composition to the vicinity of the teeth.

In general, the composition may be applied to the teeth for a period of from several
5 minutes, say from 1 minute to 5 minutes, up to one hour or so. Afterwards, the composition is preferably removed from the buccal cavity by rinsing.

Treatment by a dental surgeon will normally begin with a conventional scale-and-polish treatment. This may be followed by one or more applications of a bleaching
10 agent, eg at daily or weekly intervals, in a conventional manner. The fluorescent whitening agent may then be applied after the last bleaching treatment.

Maintenance treatment may then be performed by the patient, eg by regular use of a toothpaste containing relatively low concentrations of fluorescent whitening
15 agent and optionally a bleaching agent or other tooth whitening agent. Further clinical treatments may then be required, eg at six or twelve month intervals, to maintain the whiteness of the teeth.

The compositions according to the invention may, as appropriate, contain further
20 components which are well known to those skilled in the art of dental preparations such as, for example, humectants, surfactants, abrasives, fluoride sources, desensitising agents, flavourings, colourings, sweeteners, antimicrobial agents to act as preservatives, bactericides and/or anti-plaque agents, anti-calculus (anti-tartar) agents, structuring agents, chelating agents, whitening agents, vitamins and
25 any other therapeutic actives. The choice and amount of additives for any particular form of composition will be evident to, or readily determined by, those skilled in the art.

Suitable humectants for use in dentifrice compositions include polyhydric alcohols
30 such as xylitol, sorbitol, glycerol, propylene glycol and polyethylene glycols. Sorbitol, and mixtures of glycerol and sorbitol or sorbitol and xylitol, are particularly effective. A humectant helps to prevent dentifrice compositions from hardening on

exposure to air, and may also provide a moist feel, smooth texture, flowability, and a desirable sweetness in the mouth. Suitably, such humectants may comprise from about 0-85%, preferably from about 20-80% by weight of the composition.

- 5 Suitable surfactants for use in dentifrices, mouthwashes etc. are usually water-soluble organic compounds, and may be anionic, non-ionic, cationic or amphoteric species.

10 Anionic surfactants include the water-soluble salts of C10-C18 alkyl sulphates (eg sodium lauryl sulfates), water soluble salts of C10-C18 ethoxylated alkyl sulphates, water soluble salts of C10-C18 alkyl sarcosinates, the water-soluble salts of sulfonated monoglycerides of C10-C18 fatty acids (eg sodium coconut monoglyceride sulfonates), alkyl aryl sulfonates (eg sodium dodecyl benzene sulfonate), sodium salts of the coconut fatty acid amide of N-methyltaurine and
15 sodium salts of long chain olefin sulfonates (eg sodium C14-C16 olefin sulfonates).

20 Non-ionic surfactants suitable for use in oral compositions include the products of the condensation of alkylene oxide groups with aliphatic or alkylaromatic species, and may be for example, polyethylene oxide condensates of alkyl phenols, ethylene oxide/propylene oxide copolymers (available from BASF Wyandotte Chemical Corporation under the trade name 'Pluronic'), ethylene oxide/ethylene diamine copolymers, ethylene oxide condensates of aliphatic alcohols, long chain tertiary amine oxides, long chain tertiary phosphine oxides, long chain dialkyl
25 sulfoxides and mixtures thereof. Alternatives include ethoxylated sorbitan esters such as those available from ICI under the trade name "Tween".

30 Cationic surfactants are generally quaternary ammonium compounds having one C8-C18 alkyl chain and include, for example, lauryl trimethylammonium chloride, cetyl trimethylammonium bromide, cetyl pyridinium chloride, diisobutylphenoxyethoxyethyl dimethylbenzylammonium chloride, coconut alkyl trimethylammonium nitrite and cetyl pyridinium fluoride.

Also useful are benzyl ammonium chloride, benzyl dimethyl stearyl ammonium chloride, and tertiary amines having one C1-C18 hydrocarbon group and two (poly)oxyethylene groups.

5

Amphoteric surfactants may be aliphatic secondary and tertiary amines comprising aliphatic species which may be branched or unbranched, and in which one of the aliphatic species is a C8-C18 species and the other contains an anionic hydrophilic group, for example, sulfonate, carboxylate, sulfate, phosphonate or phosphate. Examples of quaternary ammonium compounds are the quaternized imidazole derivatives available under the trade name 'Miranol' from the Miranol Chemical Company. Other amphoteric surfactants that may be employed are fatty acid amido alkyl betaines where one alkyl group is commonly C10-C12 such as cocamido propyl betaine, for example Tego Betain supplied by T H Goldschmitt.

15

Flavouring agents may be added to increase palatability and may include, for example, menthol, oils of peppermint, spearmint, wintergreen, sassafras and clove. Sweetening agents may also be used, and these include D-tryptophan, saccharin, dextrose, aspartame, levulose, acesulfam, dihydrochalcones and sodium cyclamate..

20

Colouring agents and pigments may be added to improve the visual appeal of the composition. Suitable colourants include dyes and pigments. A suitable and commonly used pigment is pigment grade titanium dioxide, which provides a strong white colour.

25

Suitably, as described above, the compositions of the invention may include a further antimicrobial agent as a preservative, antibacterial and/or anti-plaque agent. Suitable antimicrobial agents include water soluble sources of certain metal ions such as zinc, copper and silver such as zinc citrate and silver chloride, cetyl pyridinium chloride, the bis-biguanides (such as chlorhexidine), aliphatic amines, phenolics such as bromochlorophene and triclosan, salicylanilides and

30

quaternary ammonium compounds. Optionally, the formulations may also contain enzymes that will disrupt the pellicle or interfere with bacterial intercellular polysaccharides. Examples include proteases such as papain and bromelain or dextranases. Natural enzymatic biocidal systems such as a system comprising
5 lactoperoxidase and glucose oxidase may also be employed.

The composition may also comprise an anti-calculus (anti-tartar) agent. Suitable anti-calculus agents include zinc salts such as zinc citrate and zinc chloride, polyphosphates and pyrophosphates. Suitable pyrophosphates include the
10 sodium and potassium pyrophosphates, preferably disodium pyrophosphate, dipotassium pyrophosphate, tetrasodium pyrophosphate and tetrapotassium pyrophosphate. A preferred source of pyrophosphate is a mixture of tetrasodium pyrophosphate and tetrapotassium pyrophosphate. Suitable polyphosphates include sodium tripolyphosphate.

15 Structuring (gelling) agents may be required in, for example, dentifrices and gums to provide desirable textural properties and "mouthfeel". Suitable agents include natural gum binders such as gum tragacanth, xanthan gum, gum karaya and gum arabic, seaweed derivatives such as Irish moss and alginates, smectite clays such
20 as bentonite or hectorite, carboxyvinyl polymers and water-soluble cellulose derivatives such as hydroxyethyl cellulose and sodium carboxymethyl cellulose. Improved texture may also be achieved, for example, by including colloidal magnesium aluminium silicate.

25 Suitable vitamins for inclusion in the dental preparations of the present invention include vitamins A, B5, B6, C and E.

The composition may also comprise a polymer to enhance delivery and retention of the active ingredients to the tooth surface. Suitable polymers may include
30 PVM/MA (poly(vinylmethylether/maleic acid)) copolymers eg Gantrez S-97 from ISP, PVP (polyvinylpyrrolidone) or PVP/VA (polyvinylpyrrolidone/vinyl acetate)

copolymers, eg Plasdene S-630 from ISP, and similar materials from other suppliers.

Where the composition contains any of the further ingredients or excipients
5 mentioned above, the concentrations of such ingredients may be generally similar to those conventional in dental compositions of the corresponding form.

The invention is illustrated by the following non-limiting Examples, in which

- 10 a) Examples 1 to 3 are examples of compositions containing a relatively low concentration of fluorescent whitening agent, suitable for home use;
b) Examples 4 to 6 are examples of compositions containing a relatively high concentration of a fluorescent whitening agent and, in the case of Example 6, a bleaching agent, these compositions being suitable for professional use; and
15 c) Example 7 is a toothpaste formulation suitable for home use, containing both fluorescent whitening agent and bleaching agent.

In all cases, the formulations can generally be prepared by simple admixture of the various ingredients, in a manner that will be evident to, or readily determined by, those skilled in the art.

20

Example 1

Tooth Gel

	<u>Material</u>	<u>% w/w</u>
	Sorbitol 70% solution	68.00
25	PEG 1500	1.39
	Sodium hydroxide	0.05
	Sodium monofluorophosphate	0.80
	Sodium saccharin	0.26
	Silica abrasive	6.94
30	Silica Thickener	9.17
	SLS Needles	1.00
	Flavouring	1.00

Sodium carboxymethylcellulose	1.00
Tinopal CBS-X	0.005
Purified water	q.s.

5 Example 2Toothpaste

	<u>Material</u>	<u>% w/w</u>
	Sorbitol 70% solution	50.00
	PEG 1500	1.39
10	Sodium hydroxide	0.05
	Sodium fluoride	0.22
	Sodium saccharin	0.26
	Silica abrasive	6.94
	Silica Thickener	9.17
15	SLS Needles	1.00
	Flavouring	1.00
	Sodium carboxymethylcellulose	1.00
	Tinopal CBS-X	0.005
	Titanium dioxide	0.50
20	Purified water	q.s.

Example 3Mouthwash

	<u>Ingredients</u>	<u>% w/w</u>
25	Sorbitol 70% solution	10.00
	Sodium Fluoride	0.05
	Sodium Saccharin	0.03
	Cetyl Pyridinium Chloride	0.05
	Ethanol 96%	7.00
30	Flavouring	0.15
	Tween 20	0.30
	Tinopal CBS-X	0.005

Purified Water

q.s.

Example 4Gel

5	<u>Ingredients</u>	<u>% w/w</u>
	Pluronic F-127	22.00
	Glycerin	10.00
	Sodium Saccharin	0.26
	Flavouring	1.00
10	Tinopal CBS-X	0.05
	Purified Water	q.s.

Example 5Toothpaint

15	<u>Ingredients</u>	<u>% w/w</u>
	Pluronic F-127	5.00
	Glycerin	10.00
	Ethanol 96%	30.00
	Tinopal CBS-X	0.05
20	Purified Water	q.s.

Example 6Gel with Bleach and FWA

	<u>Ingredients</u>	<u>% w/w</u>
25	Pluronic F-127	25.00
	Glycerin	35.00
	Carbamide Peroxide	15.00
	Tinopal CBS-X	0.05
	Sodium Saccharin	0.26
30	Flavouring	1.00
	Purified Water	q.s.

Example 7Toothpaste with Bleach and FWA

	<u>Ingredients</u>	<u>% w/w</u>
	Glycerin	35.00
5	Sodium Saccharin	0.26
	Sodium Fluoride	0.22
	Silica Abrasive	6.67
	Silica Thickener	10.83
	SCMC	1.00
10	Titanium Dioxide	0.50
	SLS	1.50
	PEG 1500	1.39
	Flavouring	1.00
	Carbamide Peroxide	0.30
15	Tinopal CBS-X	0.005
	Purified Water	q.s.

The efficacy of the method of the invention (employing Tinopal CBS-X, referred to as FWA, as fluorescent whitening agent and hydrogen peroxide as bleaching agent) is demonstrated by the following measurements.

Methods1) Treatment of teeth

25 Extracted caries-free teeth collected from dental hospitals were briefly brushed to remove any skin tissue attached. The teeth were then sterilized by autoclaving at 121°C for 25 mins. All teeth were stored in an airtight plastic container containing a solution of Phosphate Buffered Saline (PBS) with a few crystals of thymol as an antibacterial agent until required for use. Twenty four hours before the study,

30 each tooth was glued onto a numbered glass microscope slide. A pellicle was formed on the teeth by soaking in sterilised artificial saliva with the composition set out in Table 1.

After the 24 hr period of pellicle formation, the teeth were then brushed with an Oral B Plaque remover electric toothbrush for 2 mins, with a standard fluoride toothpaste, the composition of which is set out in Table 2:

5

Excess toothpaste was removed by briefly rinsing each tooth with deionised water. The teeth were returned to a fresh aliquot of artificial saliva and then measured. The measured whiteness index (determined by image analysis as described below) was taken as the baseline against which subsequent measurements were compared.

10

The teeth were bleached by immersion in a 15% hydrogen peroxide solution (prepared from 45g Urea Hydrogen Peroxide in 55g deionised water) for 1 hr. Teeth were returned to a fresh aliquot of artificial saliva and a Post-Bleach measurement taken.

15

FWA was applied to the teeth by immersion in a 500 ppm FWA solution (50 mg FWA in 100 ml deionised water stirred for 30 minutes) for 1 hr. The teeth were returned to a fresh aliquot of artificial saliva and a post-FWA measurement taken.

20

2) Image Analysis

Each tooth was imaged using a Kodak DCS 410 camera, which was mounted 10cm above and perpendicular to the specimen, using aperture F11 and a shutter speed of 1/10 second. The standardised lighting source comprised two UV tubes and two 6500K 'daylight' tubes. Each image was transferred to Adobe PhotoShop software. For each image, an 8mm² area was analysed and an average Adobe L,a,b reading obtained using the software. All L,a,b values were converted to CIE L*,a*,b* and then to the CIE whiteness index *W*, using the equation :

30

$$W = Y + 800(x_n - x) + 1700(y_n - y)$$

where Y is the Y-tristimulus value of the specimen, x and y , and x_n and y_n are the chromaticity coordinates for the specimen and for the perfect diffuser for the CIE 1931 2° Standard Observer, respectively.

5 Results

Six groups of three teeth were studied. For each group of teeth, treatment with 15% H_2O_2 produced an increase in whiteness index, but further treatment with FWA produced a further increase in whiteness. For the whole group of teeth studied, the average percentage increase in whiteness index was 17% for the 10 H_2O_2 alone, and 24% for the combined treatment using H_2O_2 and FWA.

Measurements using other FWAs

Similar measurements to those described above were made using the following 15 commercially-available FWAs in place of the Tinopal CBS-X:

1,8 Naphthalimide (Sigma Aldrich)

Blankophor CLE (stilbene derivative – Bayer Speciality Products)

Coumarin 102 (Sigma Aldrich)

20 Tinopal 5BM-GX (stilbene derivative – Ciba)

Tinopal AMS-GX (stilbene derivative – Ciba)

Tinopal LMS-X (stilbene derivative – Ciba)

In each case, a measurable increase in whiteness index was observed compared 25 with treatment with H_2O_2 alone.

Measurements using alternative bleaching agent

Similar measurements were made using sodium chlorite (generating chlorine 30 dioxide in use) as bleaching agent in place of H_2O_2 , and Tinopal CBS-X as FWA. For the group of teeth studied, the average percentage increase in whiteness

index was 8% for the bleach alone, and 41% for the combined treatment using chlorine dioxide and FWA.

Table 1 - Formula for Artificial Saliva

Material	Concentration
Calcium Chloride	2.0mM
Magnesium Chloride	4.2mM
Sodium Chloride	4.0mM
Sodium Thiocyanate	1.3mM
Sodium Fluoride	15 μ M
Potassium Iodide	8.0nM
Potassium Bicarbonate	5.0mM
Potassium Dihydrogen Phosphate	6.0mM
Potassium Chloride	10.0mM
Urea	3.2mM
Glucose	0.55mM
Lactoferrin	5.4 mg/l
Lysozyme	0.264 g/l
α -Amylase	0.38 g/l
Lactoperoxidase	2.0 mg/l
Serum Albumin	2.2 g/l

Table 2 - Composition of standard toothpaste

5

Material	% w/w
Sorbitol 70% soln	25.00
Sodium Saccharin	0.26
Sodium Fluoride	0.22
Silica Abrasive (Medium)	6.67
Silica Thickener	10.83
Sodium Carboxymethyl Cellulose	1.50
Titanium Dioxide	0.50
Sodium Lauryl Sulphate	1.50
Polyethylene Glycol 1500	1.39
Sodium Bicarbonate	1.00
Flavour	1.00
Sodium Benzoate	0.50
Purified Water	To 100%

Claims

1. A dental composition comprising a fluorescent whitening agent and a dentally acceptable diluent or carrier.

5


2. A dental composition as claimed in Claim 1, which is formulated as a toothpaste, mouthrinse, toothgel, tooth paint or dental gel.

3. A dental composition as claimed in Claim 1 or Claim 2, wherein the
10 fluorescent whitening agent absorbs light of wavelength less than 380nm and re-emits light in the wavelength range 400nm to 450nm.

4. A dental composition as claimed in any preceding claim, wherein the fluorescent whitening agent is selected from the group consisting of


15

a) bis-styrylbiphenyl compounds having the following chromophore system:




b) derivatives of stilbene having the following chromophore system:

20




c) coumarins having the following chromophore system:

24




d) naphthalimide derivatives having the following chromophore system:



5

e) benzoxazole derivatives having the following chromophore system:





10

in which R represents a conjugated group; and

f) pyrazoline and bis-pyrazoline derivatives having the following chromophore systems:


15



5. A dental composition as claimed in any preceding claim, wherein the fluorescent whitening agent is a bis-styrylbiphenyl compound of the general formula:

20

25




in which R_1 is $-\text{SO}_3\text{M}$ and R_2 , R_3 and R_4 , which may be the same or different, are
 5 selected from R_5 , $-\text{SO}_3\text{M}$, halogen (particularly Cl), $-\text{CN}$, $-\text{OC}(=\text{O})\text{R}_5$, $-\text{COOR}_5$, $-\text{SO}_2\text{N}(\text{R}_5)_2$ and $-\text{CON}(\text{R}_5)_2$, wherein R_5 represents hydrogen or C_{1-8} alkyl and M represents hydrogen or a Group I metal, eg Na, K or Li.


6. A dental composition as claimed in Claim 5, wherein R_3 is the same as R_1 ,
 10 and R_2 and R_4 are the same and are selected from R_5 , halogen, $-\text{CN}$, $-\text{OC}(=\text{O})\text{R}_5$, $-\text{COOR}_5$, $-\text{SO}_2\text{N}(\text{R}_5)_2$ and $-\text{CON}(\text{R}_5)_2$.

7. A dental composition as claimed in Claim 4, wherein the fluorescent
 whitening agent is selected from the group consisting of disodium 4,4'-bis[(4-
 15 anilino-6-morpholino-1,3,5-triazin-2-yl)amino]stilbene-2,2'-disulfonate, disodium
 4,4'-bis[[4-anilino-6-(N-methyl-N-2-hydroxyethyl)amino-1,3,5-triazin-2-yl]
 amino]stilbene-2,2'-disulfonate, and disodium 4,4'-bis[(4-anilino-6-methylamino-
 1,3,5-triazin-2-yl)aminolstilbene-2,2'-disulfonate.

20 8. A dental composition as claimed in Claim 4, wherein the fluorescent
 whitening agent is:



9. A dental composition as claimed in Claim 4, wherein the fluorescent whitening agent is:



5

10. A dental composition as claimed in Claim 4, wherein the fluorescent whitening agent is a bis-styrylbiphenyl compound.

11. A dental composition as claimed in Claim 10, wherein the fluorescent whitening agent is 4,4'-bis(2-sulfoxy)styrylbiphenyl or a salt or other soluble derivative thereof.

12. A dental composition as claimed in Claim 11, wherein the fluorescent whitening agent is disodium 4,4'-bis(2-sulfoxy)styrylbiphenyl.

15

13. A dental composition as claimed in any preceding claim, wherein the concentration of fluorescent whitening agent in the composition is less than 1,000 ppm.

20

14. A dental composition as claimed in Claim 13, wherein the concentration of fluorescent whitening agent in the composition is in the range 50ppm to 500ppm.

15. A dental composition as claimed in any Claim 13, wherein the concentration of fluorescent whitening agent in the composition is less than 100ppm.

25

16. A dental composition as claimed in Claim 15, wherein the concentration of fluorescent whitening agent in the composition is in the range 5ppm to 50ppm.

5 17. A dental composition as claimed in any preceding claim, wherein the composition comprises an additional tooth whitening agent in addition to the fluorescent whitening agent.

18. A dental composition as claimed in Claim 17, wherein the additional tooth whitening agent is a bleaching agent.

10

19. A dental composition as claimed in Claim 18, wherein the bleaching agent is a peroxide.

15 20. A dental composition as claimed in Claim 19, wherein the peroxide is hydrogen peroxide or a compound that generates hydrogen peroxide in use.

21. A dental composition as claimed in Claim 18, wherein the bleaching agent is a chlorite bleaching agent.

20 22. A method for improving or enhancing the appearance of the teeth, in particular for whitening of the teeth, which method comprises application to the teeth of a dental composition as claimed in any one of Claims 1 to 14.

25 23. A method as claimed in Claim 22, wherein the method further comprises the application to the teeth of an additional tooth whitening agent.

24. A method as claimed in Claim 23, wherein the additional tooth whitening agent is a bleaching agent.

28

25. A method as claimed in Claim 24, wherein the bleaching agent is a peroxide.

26. A method as claimed in Claim 25, wherein the peroxide is hydrogen
5 peroxide or a compound that generates hydrogen peroxide in use.

27. A method as claimed in any one of Claims 23 to 26, wherein application of the additional tooth whitening agent precedes application of the fluorescent whitening agent.

10

28. A method as claimed in any one of Claims 23 to 26, wherein application of the additional tooth whitening agent is simultaneous with application of the fluorescent whitening agent.

15

29. A method as claimed in any one of Claims 23 to 26, comprising a first stage in which the fluorescent whitening agent and the additional tooth whitening agent are applied, simultaneously or sequentially, by a dental surgeon, and a second stage in which the fluorescent whitening agent and the additional tooth whitening agent are applied, simultaneously or sequentially, by the patient.

20

30. A method as claimed in Claim 29, wherein in the first stage the additional tooth whitening agent is applied first, followed by the fluorescent whitening agent.

25

31. A method as claimed in Claim 29 or Claim 30, wherein in the second stage, the additional tooth whitening agent and the fluorescent whitening agent are applied simultaneously.

30

32. A dental composition substantially as hereinbefore described with reference to any of the Examples.

33. A method for improving or enhancing the appearance of the teeth, in particular for whitening of the teeth, substantially as hereinbefore described.

INTERNATIONAL SEARCH REPORT

PCT/GB 03/04162

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A61K7/16

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE WPI Section Ch, Week 197143 Derwent Publications Ltd., London, GB; Class A64, AN 1971-69239S XP002269380 & JP 46 037182 A (DAINICHIESEIKA COLOUR AND) abstract	1-4, 22, 32, 33
X	DATABASE WPI Section Ch, Week 197143 Derwent Publications Ltd., London, GB; Class A64, AN 1971-69238S XP002269381 & JP 46 037181 B (DAINICHIESEIKA COLOUR AND) abstract	1-4, 9, 22, 32, 33
	-/-	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *8* document member of the same patent family

Date of the actual completion of the international search

6 February 2004

Date of mailing of the international search report

19/02/2004

Name and mailing address of the ISA
 European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Fischer, J.P.

INTERNATIONAL SEARCH REPORT

PCT/GB 03/04162

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE WPI Section Ch, Week 197143 Derwent Publications Ltd., London, GB; Class A64, AN 1971-69237S XP002269382 & JP 46 037180 A (DAINICHISEIKA COLOR & CHEM MFG) abstract	1-4, 9, 22, 32, 33
X	DATABASE WPI Section Ch, Week 197108 Derwent Publications Ltd., London, GB; Class D18, AN 1971-15011S XP002269383 & JP 46 007389 B (DAINICHISEIKA COLOR & CHEM MFG) abstract	1-4, 22, 32, 33
X	DATABASE WPI Section Ch, Week 197007 Derwent Publications Ltd., London, GB; Class A36, AN 1970-10184R XP002269384 & JP 45 002669 B (DAIN) abstract	1-4, 22, 32, 33
X	DATABASE WPI Section Ch, Week 197143 Derwent Publications Ltd., London, GB; Class A64, AN 1971-69241S XP002269385 & JP 46 037184 B (DAINICHISEIKA COLOR & CHEM MFG) abstract	1-4, 22, 32, 33
X	DATABASE WPI Section Ch, Week 197143 Derwent Publications Ltd., London, GB; Class A64, AN 1971-69240S XP002269386 & JP 46 037183 B (DAINICHISEIKA COLOR & CHEM MFG) abstract	1-4, 22, 32, 33
X	DATABASE WPI Section Ch, Week 197143 Derwent Publications Ltd., London, GB; Class A64, AN 1971-69236S XP002269387 & JP 46 037179 B (DAINICHISEIKA COLOR & CHEM MFG) abstract	1-4, 22, 32, 33
	-/-	

INTERNATIONAL SEARCH REPORT

PCT/GB 03/04162

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE WPI Section Ch, Week 197007 Derwent Publications Ltd., London, GB; Class A36, AN 1970-10185R XP002269388 & JP 45 002670 B (DAIN) abstract</p>	1-4, 22, 32, 33
X	<p>DATABASE WPI Section Ch, Week 200304 Derwent Publications Ltd., London, GB; Class B02, AN 2003-042510 XP002269389 & JP 2002 234884 A (GIFUKEN) 23 August 2002 (2002-08-23) abstract</p>	1-7, 22, 32, 33
X	<p>DE 35 28 709 A (SCHLEICHER PETER) 19 February 1987 (1987-02-19) the whole document</p>	1-4, 13-16, 32
X	<p>US 3 223 588 A (PICKEL FRANK D) 14 December 1965 (1965-12-14) column 1, line 32 - line 45 column 2, line 7 - line 41; claims 1-4; example 1</p>	1-4, 13-16, 32
X	<p>WO 00/16737 A (MCLAUGHLIN GERALD G) 30 March 2000 (2000-03-30) the whole document</p>	1-4, 13-20, 22-26, 28, 32, 33
X	<p>WO 99/37236 A (ULTRADENT PRODUCTS INC) 29 July 1999 (1999-07-29) the whole document</p>	1-4, 13-20, 22-26, 28, 32, 33
X	<p>CH 568 349 A (CIBA GEIGY AG) 31 October 1975 (1975-10-31) the whole document</p>	1-4, 10-12, 22, 32, 33
X	<p>CH 532 629 A (CIBA GEIGY AG) 15 January 1973 (1973-01-15) the whole document</p>	1-4, 22, 32, 33
X	<p>FR 1 506 818 A (CIBA GEIGY) 22 December 1967 (1967-12-22) the whole document</p>	1-4, 32
X, P	<p>WO 02/098349 A (LIPO CHEMICALS INC) 12 December 2002 (2002-12-12) the whole document</p>	1-4, 22, 32, 33

INTERNATIONAL SEARCH REPORT

PCT/GB 03/04162

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
JP 46037182	A	NONE	
JP 46037181	B	NONE	
JP 46037180	A	NONE	
JP 46007389	B	NONE	
JP 45002669	B	NONE	
JP 46037184	B	NONE	
JP 46037183	B	NONE	
JP 46037179	B	NONE	
JP 45002670	B	NONE	
JP 2002234884	A	23-08-2002	NONE
DE 3528709	A	19-02-1987	DE 3528709 A1 19-02-1987 EP 0217041 A1 08-04-1987 JP 7076168 B 16-08-1995 JP 62039516 A 20-02-1987
US 3223588	A	14-12-1965	NONE
WO 0016737	A	30-03-2000	AU 6251899 A 10-04-2000 WO 0016737 A1 30-03-2000
WO 9937236	A	29-07-1999	US 6391283 B1 21-05-2002 AU 2336399 A 09-08-1999 WO 9937236 A1 29-07-1999 US 6387353 B1 14-05-2002
CH 568349	A	31-10-1975	CH 568349 A5 31-10-1975
CH 532629	A	15-01-1973	CH 532608 A 15-01-1973 AT 305946 B 26-03-1973 BE 760402 A1 16-06-1971 CA 983508 A1 10-02-1976 CS 154310 B2 29-03-1974 DE 2060288 A1 24-06-1971 ES 387272 A1 16-12-1975 FR 2073875 A5 01-10-1971 GB 1326230 A 08-08-1973 JP 49001275 B 12-01-1974 NL 7018353 A 21-06-1971 SU 406364 A3 05-11-1973 US 3872094 A 18-03-1975
FR 1506818	A	22-12-1967	CH 466507 A 15-12-1968 BE 692077 A 29-05-1967 GB 1134767 A 27-11-1968
WO 02098349	A	12-12-2002	US 2002192248 A1 19-12-2002 US 2002192260 A1 19-12-2002 FR 2825274 A1 06-12-2002

INTERNATIONAL SEARCH REPORT

PCT/GB 03/04162

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 02098349 A		FR 2829026 A1	07-03-2003
		WO 02098374 A1	12-12-2002
		WO 02098349 A2	12-12-2002
		US 2003104022 A1	05-06-2003
		US 2003152537 A1	14-08-2003
		US 2003147821 A1	07-08-2003
		US 2003170189 A1	11-09-2003